



SHIFT - 1

QUESTIONS & SOLUTIONS

Reproduced from Memory Retention

 30 JANUARY, 2023

 9:00 AM to 12:00 Noon

Duration : 3 Hours

Maximum Marks : 300

SUBJECT - CHEMISTRY

RESULT JEE ADVANCED 2022

Created HISTORY
in Just 3 Years of Inception

BEST RANK IN
**NORTH INDIA &
IIT DELHI ZONE**

AIR
5

**MAYANK
MOTWANI**

Roll No. : 20771637
Classroom Student
Since 2020



KVPY 2020
(Stream - SA)

AIR 1

KVPY 2021
(Stream - SX)

AIR 6

JEE MAIN
2022

AIR 15

Accomplish your **DREAM** with
Reliable **FACULTY TEAM**

TARGET JEE Adv. 2024

Not satisfied with your
JEE Performance ?

Join

VISHESH

For Class XII Passed /
Repeater Students

**One Year Classroom Course
for Complete
JEE (Main+Adv) Syllabus**

STARTING FROM :

15 & 29 MARCH'23

CHEMISTRY

1. Which of the following is water soluble?

- (a) BeSO_4 (b) MgSO_4 (c) CaSO_4 (d) SrSO_4
(e) BaSO_4

- (1) (a) only (2) (a) & (b) (3) (c) only (4) (c) & (d)

Ans. (2)

Sol. BeSO_4 & MgSO_4 are water soluble.

2. During the qualitative analysis of SO_3^{2-} using dil. H_2SO_4 , SO_2 gas is evolved which turns $\text{K}_2\text{Cr}_2\text{O}_7$ solution.

- (1) Green (2) Black (3) Blue (4) Red

Ans. (1)

Sol. $\text{SO}_3^{2-} \xrightarrow[\text{H}_2\text{SO}_4]{\text{dil}} \text{SO}_2 \xrightarrow[\text{H}^+]{\text{K}_2\text{Cr}_2\text{O}_7} \text{Cr}^{3+} + \text{SO}_4^{2-}$
(green)

3. Match the following

Atomic number			
(a)	52	(p)	s block
(b)	37	(q)	p block
(c)	65	(r)	d block
(d)	78	(s)	f block

- (1) a – (q), b – (p), c – (r), d – (s) (2) a – (q), b – (p), c – (s), d – (r)
(3) a – (s), b – (r), c – (p), d – (q) (4) a – (r), b – (p), c – (q), d – (s)

Ans. (2)

Sol. 52 → p-block
37 → s-block
65 → f-block
78 → d-block

4. If volume of ideal gas is increased isothermally than its internal energy
- (1) Increased (2) Remain constant
(3) Decreased (4) Can be increased or decreased

Ans. (2)

Sol. Isothermal process

$$\Delta T = 0$$




$$\Delta U = nC_v dT = 0$$

5. Which of the following compounds acts as an inhibitor for cancer growth.
- (1) Cisplatin (2) EDTA (3) Cobalt (4) Ethane 1,2-diamine

Ans. (1)

Sol. Cisplatin $[\text{Pt}(\text{NH}_3)_2\text{Cl}_2]$

6. Order of strength of ligands S^{2-} , CO , en, $\text{C}_2\text{O}_4^{2-}$, NH_3



Ans. (1)

Sol. Order of strength of ligand $\text{S}^{2-} < \text{C}_2\text{O}_4^{2-} < \text{NH}_3 < \text{en} < \text{CO}$


7. Number of lone pairs in central atom of following species
 IF_7 , ICl_4^- , XeF_2 & XeO_3
- (1) 0, 2, 3, 1 (2) 3, 2, 1, 0 (3) 1, 2, 0, 3 (4) 0, 2, 3, 1

Ans. (1)

Species	No. of lone pair
IF_7	0
ICl_4^-	2
XeF_2	3
XeO_3	1

8. Which of the following reaction can be used to prepared LiAlH_4
 (1) $\text{LiCl} + \text{AlCl}_3$ (2) $\text{LiH} + \text{Al}(\text{OH})_3$ (3) $\text{LiH} + \text{AlCl}_3$ (4) None of these

Ans. (3)



9. Permanganate reacts in acidic medium to produce Mn^{2+} . Calculate number of electrons used.

Ans. 5



10. Speed of e^- in 7th orbit is 3.6×10^6 m/s then find speed in 3rd orbit
 (1) 3.6×10^6 m/s (2) 8.4×10^6 m/s (3) 7.5×10^6 m/s (4) 1.8×10^6 m/s

Ans. (2)

Sol. $V = 2.18 \times 10^6 \times \frac{Z}{n}$ m/s

$$3.6 \times 10^6 = 2.18 \times 10^6 \times \frac{Z}{7} \quad \dots\dots (1)$$

$$V = 2.18 \times 10^6 \times \frac{Z}{3}$$

$$= \frac{3.6 \times 10^6}{V} = \frac{Z}{7} \times \frac{3}{Z}$$

$$= \frac{3.6 \times 10^6}{V} = \frac{1 \times 3}{7}$$

$$V = \frac{3.6 \times 10^6 \times 7}{3}$$

$$= 8.4 \times 10^6 \text{ m/s}$$

11. If rate constant K is 2.011 min^{-1} for radioactive decay reaction. Calculate time period for changing mass of radioactive element from 7 gram to 2 gram.
 [$\log_{10} 7 = 0.84$, $\log_{10} 2 = 0.30$]

Ans. 0.618 min.

Sol. $t = \frac{1}{K} \ln \left[\frac{7}{2} \right]$

$$= \frac{1}{2.011} \ln 3.5$$

$$= \frac{2.303}{2.011} \log_{10} 3.5$$

$$= \frac{2.303}{2.011} [0.84 - 0.30]$$

$$= \frac{2.303}{2.011} \times 0.54 = 0.618$$

12. Molarity of CO_2 in soft drink is 0.2M. The volume of soft drink is 300 ml. Volume of CO_2 (in L) at STP present in soft drink is

Ans. 1.362 L

Sol. $n_{\text{CO}_2} = M \times V = \frac{0.2 \times 300}{1000} = \frac{6}{100}$

$$V_{\text{CO}_2} \text{ at STP} = \frac{6}{100} \times 22.7$$

$$= 1.362 \text{ L}$$

13. Find mole of a non-volatile solute dissolved in 30g water. The solution have boiling point 373.52K & $K_b(\text{water}) = 0.52 \text{ K Kg/mol}$.

Ans. 0.03 mole

Sol. ($i = 1$) Considering solute to be non-electrolyte

$$\Delta T_b = K_b \cdot m \quad \Delta T_b = 373.52 - 373 = 0.52 \text{ K}$$

$$0.52 = 0.52 \times m$$


$$\Rightarrow m = 1$$

$$m = \frac{n_{\text{solute}}}{W_{\text{solvent(g)}}} \times 1000$$





$$1 = \frac{n_{\text{solute}}}{30} \times 1000$$

$$n_{\text{solute}} = 0.03 \text{ mol}$$

14. Observe the following reactions



A and B are respectively.



Ans. (2)

15. Which of the following acts as antacid?

(1) Brompheniramine

(2) Terfenadine

(3) Ranitidine

(4) Iproniazid

Ans. (3)

16. Caprolactum when heated at high temperature gives

(1) Nylon-6,6

(2) Nylon-6



(3) Teflon

(4) Buna-S

Ans. (2)

17. Match the following

Column – I



Column – II

(i) Swart's reaction

(ii) Finkelstein reaction


(iii) Fittig reaction

(iv) Sandmeyer's reaction

	p	q	r	s
(1)	iii	i	ii	iv
(2)	ii	iii	i	iv
(3)	iv	iii	ii	i
(4)	i	ii	iii	iv

Ans. (2)



18. Which of the following is correct acidic strength order for the marked hydrogen in the given compound ?



- (1) $a > d > b > c$ (2) $a > b > d > c$ (3) $c > d > b > a$ (4) $a > c > b > d$

Ans. (1)

19. Consider the following reactions.







Find A, B & C respectively

- (1) NO, O₃, O (2) O, NO, O₃
(3) NO, O, O₃ (4) O₃, O, NO

Ans. (2)

20. Which of the following compound gives positive test with Fehling solution and blood red colour when fused with sodium metal followed by neutral FeCl₃ solution?







Ans. (1)

Sol. -CHO group gives positive Tollen's test where as,



21. Which of the following reaction will yield benzyl isocyanide as a major product?



Ans. (2)


22. Mark correct answer on the basis of following two statements.

Statement-I : Ketoses give seliwanoff's test faster than aldose.

Statement-II : When heated, fructose (ketose sugar) is more rapidly dehydrated than glucose (aldose sugar).


- (1) Both statements are true.
 (2) Both statements are false.
 (3) (I) is true (II) is false.
 (4) (II) is true (I) is false.

Ans. (1)



Find the number of π bonds present in compound A

Ans. 4



Since compound gives FeCl₃ test so phenolic group is present.

Compound gives NaOI test (Iodoform test), so methyl ketone group should present.

#IITkipooritaiyyari



ADMISSIONS OPEN

(Session 2023-24)

JEE (Main + Adv.) | JEE (Main)
Junior Division (VI to X)

Appear in ONLINE

Reliable National Entrance Test (R-NET)

Test on Every

SUNDAY

Scholarship up to 90%



AIR
5

MAYANK MOTWANI
Classroom



RELIABLE INSTITUTE : A-10, Road No.1, IPIA, Kota-324005 (Rajasthan), India

Tel. : 0744-3535544, 2665544 | Website : www.reliablekota.com | E-mail : info@reliablekota.com

[f](#) reliablekota [t](#) reliablekota [y](#) reliableinstitutekota [i](#) reliable_kota [in](#) reliablekota