

## FINAL NATIONAL STANDARD EXAMINATION - 2019

(Held On Sunday 24<sup>th</sup> November, 2019)

### PHYSICS

### TEST PAPER WITH ANSWER

1. A pendulum is made by using a thread of length 300 cm and a small spherical bob of mass 100 g. It is suspended from a point S. The bob is pulled from its position of rest at O to the point A so that the linear amplitude is 25 cm. The angular amplitude in radian and the potential energy of the bob in joule at A are respectively
- (a) 0.10 and 0.10      (b) 0.083 and 0.01      (c) 0.251 and 2.94      (d) 0.083 and 0.24

**Ans. (b)**

2. Consider the following physical expressions

(I)  $\rho v^2$  ( $\rho$  : density,  $v$  : velocity)

(II)  $\frac{Y\Delta L}{L}$  ( $Y$  : Young's modulus,  $L$  : length)

(III)  $\frac{\sigma^2}{\epsilon_0}$  ( $\sigma$  : surface density of charge)

(IV)  $h\rho r g$  ( $h$  : rise of a liquid in a capillary tube of radius  $r$ )

(a) I and II only      (b) II and III only      (c) II, III and IV only      (d) I, II and III only

**Ans. (d)**

3. Two simple pendulums of lengths 1.44 m and 1.0 m start swinging together in the same phase. The two will be in phase again after a time of
- (a) 6 second      (b) 9 second      (c) 12 second      (d) 25 second

**Ans. (c)**

4. A home aquarium partly filled with water slides down an inclined plane of inclination angle  $\theta$  with respect to the horizontal. The surface of water in the aquarium

(a) remains horizontal

(b) remains parallel to the plane of the incline

(c) forms an angle  $\alpha$  with the horizon where  $0 < \alpha < \theta$

(d) forms an angle  $\alpha$  with the horizon, where  $\theta < \alpha < 90$

**Ans. (b or c)**

5. A sound source of constant frequency travels with a constant velocity past an observer. When it crosses the observer the sound frequency sensed by the observer changes from 449 Hz to 422 Hz. If the velocity of sound is 340 m/s, the velocity of the source of sound is

(a) 8.5 m/s

(b) 10.5 m/s

(c) 12.5 m/s


(d) 14.5 m/s

**Ans. (b)**





15. In an ink-jet printer, an droplet of mass  $m$  is given a negative charge  $q$  by a computer-controlled charging unit. The charged droplet then enters the region between two deflecting parallel plates of length  $L$  separated by distance  $d$  (see figure below) with a speed  $v$ . All over this region there exists a uniform downward electric field  $E$  (in the plane of paper). Neglecting the gravitational force on the droplet, the maximum charge that can be given to this droplet, so that it does not hit any of the plates, is




- (a)  $\frac{mv^2L}{Ed^2}$       (b)  $\frac{mv^2d}{EL^2}$       (c)  $\frac{md}{Ev^2L^2}$       (d)  $\frac{mv^2L^2}{Ed}$

Ans. (b)

16. A converging beam of light is pointing to P. Two observations are made with (i) a convex lens of focal length 20 cm and, (ii) a concave lens of focal length 16 cm placed in the path of the convergent beam at a distance 12 cm before the point P. It is observed that
- (a) in both cases the images are real  
 (b) in both cases the images are virtual  
 (c) for (i) the image is real and for (ii) the image is virtual  
 (d) for (i) the image is virtual and for (ii) the image is real

Ans. (a)

17. The log-log graph for a non-linear oscillator is shown below. Assuming the constants to have appropriate dimensions the relationship between time period ( $T$ ) and the amplitude ( $A$ ) can be expressed as :-



- (a)  $T = 1000 A^2$       (b)  $T = 4A^{1/2}$       (c)  $T = 4A^2 + B$       (d)  $T = 8A^3$

Ans. (a)

18. In many situations the point source emitting a wave strats moving, through the medium, with velocity  $V$  greater than the wave velocity in that medium. In such a case when source velocity ( $V$ )  $>$  wave velocity ( $v$ ), the wave front changes

- (a) from spherical to plane  
(b) from spherical to conical  
(c) from plane to spherical  
(d) from cylindrical to spherical

Ans. (b)

19. If the average mass of a smoke particle in an Indian kitchen is  $3 \times 10^{-17}$  kg, the rms speed of the smoke particles at  $27^\circ\text{C}$  is approximately :

- (a) 2 cm/sec  
(b) 2 m/sec  
(c) 2 km/sec  
(d) none of these

Ans. (a)

20. Two wires, made of same material, one thick and the other thin are joined to form one composite wire. The composite wire is subjected to the same tension throughout. A wave travels along the wire and passes the point where the two wires are joined. The quantity which changes at the joint are

- (a) frequency only  
(b) propagation speed only  
(c) wavelength only  
(d) both propagation speed and wavelength

Ans. (d)

21. The frequency of the third overtone of a closed end organ pipe equals the frequency of the fifth harmonic of an open end organ pipe. Ignoring end correction, the ratio of their lengths  $l_{\text{open}} : l_{\text{close}}$  is

- (a) 10 : 7  
(b) 10 : 9  
(c) 2 : 1  
(d) 7 : 10

Ans. (a)

22. Light of wavelength 640 nm falls on a plane diffraction grating with 12000 lines per inch. In the diffraction pattern on a screen kept at a distance of 12 cm from the grating, the distance of the second order maximum from the central maximum is

- (a) 1.81 cm  
(b) 2.41 cm  
(c) 3.62 cm  
(d) 7.25 cm


Ans. (d)

23. If the force acting on a body is inversely proportional to its speed, the kinetic energy of the body varies with time  $t$  as

- (a)  $t^0$   
(b)  $t^1$   
(c)  $t^2$   
(d)  $t^{-1}$

Ans. (b)

24. As shown in the figure, a block of mass  $m$  is hung from the ceiling by the system of springs consisting of two layers. The force constant of each of the springs is  $k$ . The frequency of vertical oscillations of the block is



- (a)  $\frac{1}{2\pi} \sqrt{\frac{k}{5m}}$       (b)  $\frac{1}{2\pi} \sqrt{\frac{4k}{5m}}$       (c)  $\frac{1}{2\pi} \sqrt{\frac{5k}{6m}}$       (d)  $\frac{1}{2\pi} \sqrt{\frac{6k}{5m}}$

Ans. (d)

25. Two simple harmonic motions are given by  $x_1 = a \sin \omega t + a \cos \omega t$  and  $x_2 = a \sin \omega t + \frac{a}{\sqrt{3}} \cos \omega t$ .

The ratio of the amplitudes of the first to the second and the phase difference between them respectively are

- (a)  $\sqrt{\frac{3}{2}}$  and  $\frac{\pi}{12}$       (b)  $\frac{\sqrt{3}}{2}$  and  $\frac{\pi}{12}$       (c)  $\frac{2}{\sqrt{3}}$  and  $\frac{\pi}{12}$       (d)  $\sqrt{\frac{3}{2}}$  and  $\frac{\pi}{6}$

Ans. (a)

26. A particle is projected from the ground with a velocity  $\vec{v} = (3\hat{i} + 10\hat{j})\text{ms}^{-1}$ . The maximum height attained and the range of the particle are respectively given by (use  $g = 10 \text{ m/s}^2$ )

- (a) 5m and 6m      (b) 3m and 10m      (c) 6m and 5m      (d) 3m and 5m


Ans. (a)

27. A 20 cm long capillary tube stands vertically with lower end just in water. Water rises up to 5 cm. If the entire system is now kept on a freely falling platform, the length of the water column in the capillary tube will be

- (a) 5 cm      (b) 10 cm      (c) Zero      (d) 20 cm

Ans. (d)

28. Position-time graph of a particle moving in a potential field is shown beside. If the mass of the particle is 1 kg its total energy is approximately



- (a)  $15.45 \times 10^{-4}$  J      (b)  $30.78 \times 10^{-4}$  J      (c)  $7.71 \times 10^{-4}$  J      (d)  $3.85 \times 10^{-4}$  J

Ans. (c)

29. An observer stands on the platform at the front edge of the first bogie of a stationary train. The train starts moving with uniform acceleration and the first bogie takes 5 seconds to cross the observer. If all the bogies of the train are of equal length and the gap between them is negligible, the time taken by the tenth bogie to cross the observer is

- (a) 1.07 s      (b) 0.98 s      (c) 0.91 s      (d) 0.81 s

Ans. (d)

30. The resistive force on an aeroplane flying in a horizontal plane is given by  $F_f = kv^2$ , where  $k$  is constant and  $v$  is the speed of the aeroplane. When the power output from the engine is  $P_0$ , the plane flies at a speed  $v_0$ . If the power output of the engine is doubled the aeroplane will fly at a speed of

- (a)  $1.12 v_0$       (b)  $1.26 v_0$       (c)  $1.41 v_0$       (d)  $2.82 v_0$

Ans. (b)

31. A 3.0 cm thick layer of oil (density  $\rho_{oil} = 800 \text{ kg/m}^3$ ) floats on water (density  $\rho_w = 1000 \text{ kg/m}^3$ ) in a transparent glass beaker. A solid cylinder is observed floating vertically with  $\frac{1}{3}$  of it in water and

$\frac{1}{3}$  in the oil. Oil is gently poured into the beaker until the cylinder floats in oil only. The fraction of the solid cylinder in oil now is

- (a)  $\frac{3}{5}$       (b)  $\frac{2}{3}$       (c)  $\frac{3}{4}$       (d)  $\frac{8}{9}$

Ans. (c)

32. A wedge of mass  $M$  rests on a horizontal frictionless surface. A block of mass  $m$  starts sliding down the rough inclined surface of the wedge to its bottom. During the course of motion, the centre of mass of the block and the wedge system
- (a) does not move at all (b) moves horizontally with constant speed  
(c) moves horizontally with increasing speed (d) moves vertically with increasing speed

Ans. (d)

33. A uniform circular disc rotating at a fixed angular velocity  $\omega$  about an axis normal to its plane and passing through its centre has kinetic energy  $E$ . If the same disc rotates with an angular velocity  $2\omega$  about a parallel axis passing through the edge, its kinetic energy will be
- (a)  $2 E$  (b)  $4 E$  (c)  $10 E$  (d)  $12 E$

Ans. (d)

34. Avalanche breakdown in a p-n junction primarily depends on the phenomenon of
- (a) doping (b) collision  
(c) recombination (d) ionization

Ans. (b)

35. A source emits photons of energy  $5 \text{ eV}$  which are incident on a metallic sphere of work function  $3.0 \text{ eV}$ . The radius of the sphere is  $r = 8 \times 10^{-3} \text{ m}$ . It is observed that after some time emission of photoelectrons from the metallic sphere is stopped. Charge on the sphere when the photoemission stops is.
- (a)  $1.77 \times 10^{-16} \text{ C}$  (b)  $1.77 \times 10^{-12} \text{ C}$   
(c)  $1.11 \times 10^{-12} \text{ C}$  (d)  $1.11 \times 10^{-10} \text{ C}$

Ans. (b)

36. The dc component of current in the output of a half - wave rectifier with peak value  $I_0$  is
- (a) zero (b)  $\frac{I_0}{\pi}$  (c)  $\frac{I_0}{2\pi}$  (d)  $\frac{2I_0}{\pi}$

Ans. (b)


37. In an experiment on photoelectric effect, the slope of straight line graph between the stopping potential and the frequency of incident radiation gives.
- (a) Electron charge(e) (b) Planck constant (h)  
(c)  $\frac{h}{e}$  (d) Work function (W)

Ans. (c)





43. In the circuit shown beside the charge on each capacitor is



(a)  $(C_1 + C_2)(E_1 - E_2)$

(b)  $\frac{C_1 C_2}{C_1 + C_2}(E_1 + E_2)$

(c)  $\frac{C_1 C_2}{C_1 + C_2}(E_1 - E_2)$

(d)  $(C_1 - C_2)(E_1 + E_2)$

Ans. (c)

44. A stationary hydrogen atom emits photon corresponding to the first line (highest wave length) of Lyman series. If  $R$  is the Rydberg constant and  $M$  is the mass of the atom, the recoil velocity of the atom is

(a)  $\frac{Rh}{4M}$

(b)  $\frac{3Rh}{M}$

(c)  $\frac{3Rh}{4M}$

(d)  $\frac{Rh}{M}$

Ans. (c)

45. Heat is absorbed or evolved when current flows in a conductor having a temperature gradient. This phenomenon is known as

(a) Joule effect

(b) Peltier effect

(c) Seebeck effect

(d) Thomson effect

Ans. (d)

46. A concave mirror has a radius of curvature  $R$  and forms the image of an object placed at a distance  $1.5R$  from the pole of the mirror. An opaque disc of diameter half the aperture of the mirror is placed with the pole at the centre. As a result :

(a) the position of the image will be the same but its central half will disappear.

(b) the position of the image will be the same but its outer half will disappear.

(c) the complete image will be seen at the same position and it will be exactly identical with the initial image.

(d) the complete image will be seen at the same position but it will not be identical in all respect with the initial image.

Ans. (d)

47. A ray of white light is made incident on the refracting surface of a prism such that after refraction at this surface, the green component falls on the second surface at its critical angle. The colours present in the emergent beam will be :

- (a) violet, indigo and blue (b) violet, indigo, blue, yellow, orange and red  
(c) yellow, orange and red (d) all colours

Ans. (c)

48. In a compound microscope, having tube-length 30 cm, the power of the objective and the eye-piece are 100 D and 10D respectively. Then the magnification produced by the microscope when the final image is at the least distance of distinct vision (25 cm) will be :

- (a) 55 (b) 64 (c) 77 (d) 90

Ans. (c)

49. Parallel rays are incident on a glass sphere of diameter 10 cm and having refractive index 1.5. The sphere converges these rays at a certain point. The distance of this point from the centre of the sphere will be :

- (a) 2.5 cm (b) 5 cm (c) 7.5 cm (d) 12.5 cm

Ans. (c)

50. A jet of water from 15 cm diameter nozzle of a fire hose can reach the maximum height of 25 m. The force exerted by the water jet on the hose is :

- (a) 4.24 kN (b) 17.32 kN (c) 2.17 kN (d) 8.66 kN


Ans. (d)

51. In an electromagnetic wave the phase difference between electric vector and magnetic vector is

- (a) Zero (b)  $\frac{\pi}{2}$  (c)  $\pi$  (d)  $\frac{3\pi}{2}$

Ans. (a)

52. Plots of intensity (I) of radiation emitted by a black body versus wavelength ( $\lambda$ ) at three different temperatures  $T_1$ ,  $T_2$  and  $T_3$  respectively are shown in figure. Choose the correct statement.



- (a)  $T_1 > T_2 > T_3$  necessarily (b)  $T_3 > T_2 > T_1$  necessarily  
(c)  $T_2 = (T_1 + T_3) / 2$  necessarily (d)  $T_2^2 = T_1 T_3$  necessarily

Ans. (a)

53. Consider a composite slab consisting of two different materials having equal thickness and equal area of cross section. The thermal conductivities are  $K$  and  $2K$  respectively. The equivalent thermal conductivity of the composite slab is :

- (a)  $\frac{2K}{3}$                       (b)  $\sqrt{2}K$                       (c)  $3K$                       (d)  $\frac{4K}{3}$

Ans. (d)

54. A large horizontal uniform disc can rotate freely about a rigid vertical axis passing through its centre  $O$ . A man stands at rest the edge of the disc at a point  $A$ . The mass of the disc is 22 times the mass of the man. The man starts moving along the edge of the disc. When he reaches  $A$ , after completing one rotation relative to the disc, the disc has turned through :

- (a)  $30^\circ$                       (b)  $90^\circ$                       (c)  $60^\circ$                       (d)  $45^\circ$

Ans. (a)

55. Two factories are sounding their sirens at 400 Hz each. A man walks from one factory towards the other at a speed of 2 m/s. The velocity of sound is 320 m/s. The number of beats heard by the person in one second will be

- (a) 6                      (b) 5                      (c) 4                      (d) 2.5

Ans. (b)

56. The temperature of an isolated black body falls from  $T_1$  to  $T_2$  in time  $t$ . Then,  $t = Cx$  where  $x$  is :

- (a)  $\left(\frac{1}{T_2} - \frac{1}{T_1}\right)$                       (b)  $\left(\frac{1}{T_2^2} - \frac{1}{T_1^2}\right)$                       (c)  $\left(\frac{1}{T_2^3} - \frac{1}{T_1^3}\right)$                       (d)  $\left(\frac{1}{T_2^4} - \frac{1}{T_1^4}\right)$

Ans. (c)

57. Two charges  $-q$  and  $-q$  are placed at points  $(0, d)$  and  $(0, -d)$ . A charge  $+q$ , free to move along  $X$  axis, will oscillate with a force proportional to

- (a)  $\frac{1}{x^2 + d^2}$                       (b)  $\frac{1}{x^2}$                       (c)  $\frac{x}{(d^2 + x^2)^{\frac{3}{2}}}$                       (d)  $\frac{1}{\sqrt{x^2 + d^2}}$


Ans. (c)

58. The average translational kinetic energy of oxygen ( $M = 32$ ) molecules at a certain temperature is 0.048 eV. The translational kinetic energy of nitrogen ( $M = 28$ ) molecules at the same temperature is (consider the two gases to be ideal) :

- (a) 0.0015 eV                      (b) 0.042 eV                      (c) 0.048 eV                      (d) 0.768 eV

Ans. (c)

59. In the following figures the velocity-time graphs for three particles 1, 2 and 3 are shown -




The magnitude of average acceleration of the three particles, over 10 s, bear the relationship


- (a)  $a_1 > a_2 > a_3$       (b)  $a_2 > a_1 > a_3$       (c)  $a_3 > a_2 > a_1$       (d)  $a_1 = a_2 = a_3$

Ans. (d)

60. The potential energy ( $U$ ) of a particle moving in a potential field varies with its displacement ( $x$ ) as shown below -



The variation of force  $F(x)$  acting on the particle as a function of  $x$  can be represented by



- (a) Fig (i)      (b) Fig (ii)      (c) Fig (iii)      (d) Fig (iv)

Ans. (d)



65. A pin of small length 'a' is placed along the axis of a concave mirror of focal length  $f$ , at the distance  $u$  ( $>f$ ) from its pole. The length of its image is 'b'. If the same object is placed perpendicular to its axis at the same distance  $u$  and the length of its image is now 'c', then :-

(a)  $b = a \frac{f^2}{(u-f)^2}$       (b)  $c = \sqrt{ab}$       (c)  $c = b \frac{u-f}{f}$       (d)  $bc = \frac{a^2 f^3}{(u-f)^3}$

Ans. (a,b,c,d)

66. A thin rod of length 10 cm is placed along the axis of a concave mirror of focal length 30 cm in such a way that one end of the image coincides with one end of the object. The length of the image may be :-

(a) 7.5 cm      (b) 12 cm      (c) 15 cm      (d) 10 cm

Ans. (a,c)

67. The mass of an electron can be expressed as :-

(a) 0.512 MeV      (b)  $8.19 \times 10^{-14} \text{ J}/c^2$       (c)  $9.1 \times 10^{-31} \text{ kg}$       (d) 0.00055 amu  
where  $c$  is speed of light in vacuum

Ans. (b,c,d)

68. Select the correct statement(s), out of the following, about diffraction at  $N$  parallel slits :-

- (a) There are  $(N-1)$  minima between each pair of principal maxima.  
(b) There are  $(N-2)$  secondary maxima between each pair of principal maxima.  
(c) Width of principal maximum is proportional to  $1/N$ .  
(d) The intensity at the principal maxima varies as  $N^2$ .


Ans. (a,b,c,d)

69. An electric dipole placed in a non-uniform electric field may experience :-

- (a) no net force, no torque      (b) a net force, but no torque.  
(c) no net force, but a torque      (d) a net force and a torque.

Ans. (a,b,c,d)

70. Two long parallel wires carry currents of equal magnitude ( $I$ ) but in opposite directions. These wires are suspended from fixed rod PQ by four chords of equal length  $L$  as shown. The mass per unit length of each wire is  $\lambda$ , the value of angle  $\theta$  subtended by two chords OA and OB, assuming it to be small, is :-



(a)  $\theta = I \sqrt{\frac{\mu_0 \lambda}{4\pi g L}}$       (b)  $\theta = I \sqrt{\frac{\mu_0}{\pi \lambda g L}}$       (c)  $\theta = I \sqrt{\frac{\mu_0 g}{4\pi \lambda L}}$       (d)  $\theta = I \sqrt{\frac{\mu_0 \lambda g}{\pi L}}$

Ans. (b)