

FINAL JEE–MAIN EXAMINATION – SEPTEMBER, 2020


(Held On Sunday 06th SEPTEMBER, 2020) TIME : 3 PM to 6 PM

CHEMISTRY

1. The value of K_C is 64 at 800 K for the reaction

$$\text{N}_2(\text{g}) + 3\text{H}_2(\text{g}) \rightleftharpoons 2\text{NH}_3(\text{g})$$

The value of K_C for the following reaction is :



- (1) $\frac{1}{4}$ (2) $\frac{1}{8}$ (3) 8 (4) $\frac{1}{64}$

Official Ans. by NTA (2)

2. The element that can be refined by distillation is :

- (1) nickel (2) zinc
 (3) gallium (4) tin

Official Ans. by NTA (2)

3. The correct match between **Item-I** and **Item-II** :

Item-I	Item-II
(a) Natural rubber	(I) 1, 3-butadiene + styrene
(b) Neoprene	(II) 1, 3-butadiene + acrylonitrile
(c) Buna-N	(III) Chloroprene
(d) Buna-S	(IV) Isoprene

- (1) (a) - (III), (b) - (IV), (c) - (I), (d) - (II)
 (2) (a) - (IV), (b) - (III), (c) - (II), (d) - (I)
 (3) (a) - (IV), (b) - (III), (c) - (I), (d) - (II)
 (4) (a) - (III), (b) - (IV), (c) - (II), (d) - (I)

Official Ans. by NTA (2)

4. Mischmetal is an alloy consisting mainly of:

- (1) lanthanoid metals
 (2) actinoid metals
 (3) actinoid and transition metals
 (4) lanthanoid and actinoid metals

Official Ans. by NTA (1)


TEST PAPER WITH ANSWER

5. Reaction of an inorganic sulphite X with dilute H_2SO_4 generates compound Y. Reaction of Y with NaOH gives X. Further, the reaction of X with Y and water affords compound Z. Y and Z, respectively, are:

- (1) S and Na_2SO_3
 (2) SO_2 and NaHSO_3
 (3) SO_3 and NaHSO_3
 (4) SO_2 and Na_2SO_3

Official Ans. by NTA (2)

6. The IUPAC name of the following compound is :



- (1) 3-amino-4-hydroxymethyl-5-nitrobenzaldehyde
 (2) 2-nitro-4-hydroxymethyl-5-aminobenzaldehyde
 (3) 4-amino-2-formyl-5-hydroxymethylnitrobenzene
 (4) 5-amino-4-hydroxymethyl-2-nitrobenzaldehyde

Official Ans. by NTA (4)

7. Dihydrogen of high purity (> 99.95%) is obtained through:

- (1) the electrolysis of warm $\text{Ba}(\text{OH})_2$ solution using Ni electrodes.
 (2) the reaction of Zn with dilute HCl
 (3) the electrolysis of brine solution.
 (4) the electrolysis of acidified water using Pt electrodes.

Official Ans. by NTA (1)

8. Match the following :

Test/Method	Reagent
(i) Lucas Test	(a) $C_6H_5SO_2Cl/aq. KOH$
(ii) Dumas method	(b) $HNO_3/AgNO_3$
(iii) Kjeldahl's method	(c) CuO/CO_2
(iv) Hinsberg Test	(d) Conc. HCl and $ZnCl_2$
	(e) H_2SO_4

- (1) (i)-(d), (ii)-(c), (iii)-(e), (iv)-(a)
 (2) (i)-(b), (ii)-(d), (iii)-(e), (iv)-(a)
 (3) (i)-(d), (ii)-(c), (iii)-(b), (iv)-(e)
 (4) (i)-(b), (ii)-(a), (iii)-(c), (iv)-(d)

Official Ans. by NTA (1)

9. The reaction of NO with N_2O_4 at 250 K gives :

- (1) N_2O_5 (2) NO_2
 (3) N_2O (4) N_2O_3

Official Ans. by NTA (4)

10. For the given cell ;

$Cu(s)|Cu^{2+}(C_1M)||Cu^{2+}(C_2M)|Cu(s)$ change in Gibbs energy (ΔG) is negative, if :

- (1) $C_1 = 2C_2$ (2) $C_2 = \frac{C_1}{\sqrt{2}}$
 (3) $C_1 = C_2$ (4) $C_2 = \sqrt{2}C_1$

Official Ans. by NTA (4)

11. A crystal is made up of metal ions ' M_1 ' and ' M_2 ' and oxide ions. Oxide ions form a ccp lattice structure. The cation ' M_1 ' occupies 50% of octahedral voids and the cation ' M_2 ' occupies 12.5% of tetrahedral voids of oxide lattice. The oxidation numbers of ' M_1 ' and ' M_2 ' are, respectively :

- (1) +2, +4 (2) +3, +1
 (3) +1, +3 (4) +4, +2




Official Ans. by NTA (1)

12. For a d^4 metal ion in an octahedral field, the correct electronic configuration is :

- (1) $t_{2g}^4 e_g^0$ when $\Delta_0 < P$
 (2) $e_g^2 t_{2g}^2$ when $\Delta_0 < P$
 (3) $t_{2g}^3 e_g^1$ when $\Delta_0 < P$
 (4) $t_{2g}^3 e_g^1$ when $\Delta_0 > P$

Official Ans. by NTA (3)

13. Which of the following compounds can be prepared in good yield by Gabriel phthalimide synthesis?



Official Ans. by NTA (1)

14. The correct match between **Item-I** (starting material) and **Item-II** (reagent) for the preparation of benzaldehyde is :

Item-I	Item-II
(I) Benzene	(P) HCl and $SnCl_2, H_3O^+$
(II) Benzonitrile	(Q) $H_2, Pd-BaSO_4, S$ and quinoline
(III) Benzoyl Chloride	(R) CO, HCl and $AlCl_3$
(1) (I)-(Q), (II)-(R) and (III)-(P)	
(2) (I)-(R), (II)-(Q) and (III)-(P)	
(3) (I)-(R), (II)-(P) and (III)-(Q)	
(4) (I)-(P), (II)-(Q) and (III)-(R)	

Official Ans. by NTA (3)

15. The average molar mass of chlorine is 35.5 g mol^{-1} . The ratio of ^{35}Cl to ^{37}Cl in naturally occurring chlorine is close to :

- (1) 4 : 1
- (2) 1 : 1
- (3) 2 : 1
- (4) 3 : 1

Official Ans. by NTA (4)

16. Which one of the following statements not true ?

- (1) Lactose contains α -glycosidic linkage between C_1 of galactose and C_4 of glucose.
- (2) Lactose ($\text{C}_{11}\text{H}_{22}\text{O}_{11}$) is a disaccharide and it contains 8 hydroxyl groups.
- (3) On acid hydrolysis, lactose gives one molecule of D(+)-glucose and one molecule of D(+)-galactose.
- (4) Lactose is a reducing sugar and it gives Fehling's test.


Official Ans. by NTA (1)

17. A set of solutions is prepared using 180 g of water as a solvent and 10 g of different non-volatile solutes A, B and C. The relative lowering of vapour pressure in the presence of these solutes are in the order [Given, molar mass of A = 100 g mol^{-1} ; B = 200 g mol^{-1} ; C = $10,000 \text{ g mol}^{-1}$]

- (1) $A > B > C$
- (2) $A > C > B$
- (3) $C > B > A$
- (4) $B > C > A$

Official Ans. by NTA (1)

18. For a reaction,



the free energy change is plotted as a function of temperature. The temperature below which the oxide is stable could be inferred from the plot as the point at which :

- (1) the slope changes from positive to zero
- (2) the free energy change shows a change from negative to positive value
- (3) the slope changes from negative to positive
- (4) the slope changes from positive to negative

Official Ans. by NTA (2)


19. Match the following compounds (Column-I) with their uses (Column-II) :

S.No.	Column – I	S.No.	Column – II
(I)	$\text{Ca}(\text{OH})_2$	(A)	casts of statues
(II)	NaCl	(B)	white wash
(III)	$\text{CaSO}_4 \cdot \frac{1}{2}\text{H}_2\text{O}$	(C)	antacid
(IV)	CaCO_3	(D)	washing soda preparation

- (1) (I)-(D), (II)-(A), (III)-(C), (IV)-(B)
- (2) (I)-(B), (II)-(C), (III)-(D), (IV)-(A)
- (3) (I)-(C), (II)-(D), (III)-(B), (IV)-(A)
- (4) (I)-(B), (II)-(D), (III)-(A), (IV)-(C)

Official Ans. by NTA (4)

20. The increasing order of the boiling points of the major products A, B and C of the following reactions will be :



- (1) $C < A < B$
- (2) $B < C < A$
- (3) $A < B < C$
- (4) $A < C < B$

Official Ans. by NTA (69)

21. For Freundlich adsorption isotherm, a plot of $\log(x/m)$ (y-axis) and $\log p$ (x-axis) gives a straight line. The intercept and slope for the line is 0.4771 and 2, respectively. The mass of gas, adsorbed per gram of adsorbent if the initial pressure is 0.04 atm, is $______ \times 10^{-4}g$.
($\log 3 = 0.4771$)
Official Ans. by NTA (48.00)
22. A solution of phenol in chloroform when treated with aqueous NaOH gives compound P as a major product. The mass percentage of carbon in P is $______$. (to the nearest integer)
(Atomic mass : C = 12; H = 1; O = 16)
Official Ans. by NTA (69.00)
Official Ans. by ALLEN (68.85)
23. If the solubility product of AB_2 is $3.20 \times 10^{-11} M^3$, then the solubility of AB_2 in pure water is $______ \times 10^{-4} mol L^{-1}$. [Assuming that neither kind of ion reacts with water]
Official Ans. by NTA (2.00)
24. The rate of a reaction decreased by 3.555 times when the temperature was changed from $40^\circ C$ to $30^\circ C$. The activation energy (in $kJ mol^{-1}$) of the reaction is $______$.
Take; $R=8.314 J mol^{-1} K^{-1}$ $\ln 3.555 = 1.268$
Official Ans. by NTA (100.00)
Official Ans. by ALLEN (99.98)
25. The atomic number of Unnilunium is $______$.
Official Ans. by NTA (101.00)